

The focus of my research in mathematical logic is definable Skolem functions in weakly o-minimal structures, specifically constructive definitions of Skolem functions. I have found an explicit construction of Skolem functions for a subclass of valuationsal weakly o-minimal theories called  $T$ -immune, where it was known that there were definable Skolem functions, but for which there was no construction. I have also analyzed a certain subclass of nonvaluational weakly o-minimal structures, regarded colloquially to be as close as possible to being o-minimal, and found that such theories in fact do not have definable Skolem functions.

I currently have a paper in preparation which narrates the results outlined above and some extensions to more general cases.

## Background

For the purpose of this introduction, a *model*  $\mathcal{M}$  is a set (called the *universe* of the model) together with a specified algebraic structure of *definable sets*, which may be subsets of the universe  $\mathcal{M}$  itself, or of  $\mathcal{M}^n$  for a finite integer  $n$ . A canonical example of a model is the field of real numbers,  $(\mathbb{R}, +, \cdot, 0, 1, <)$ , in which case the definable subsets are precisely the semialgebraic subsets of  $\mathbb{R}^n$ .

An *o-minimal structure* is a model  $\mathcal{M}$  whose definable sets include a (usually dense) linear order on the universe, and for which any definable set of  $\mathcal{M}$  is a finite union of points and intervals (whose infima and suprema are elements of the universe). The real field is the archetype for this study. During the past several decades, a rash of work has led to a powerful structure theory for general o-minimal structures (*cf.* [6] [8] [11] [12]). Among the properties enjoyed by this class, every o-minimal structure has a strong *cellular decomposition* property which guarantees all definable subsets  $\mathcal{M}^n$  can be written as a finite union of simple definable subsets, called *cells*.

Independently of this, *Skolem functions* were developed initially in order to prove what is now known as the Löwenheim-Skolem theorem (*cf.* [2]). Given a model  $\mathcal{M}$  and definable set  $D \subseteq \mathcal{M}^n$ , a Skolem function is a function  $f$  such that for every  $\bar{a} \in \mathcal{M}^{n-1}$ , if there is some  $y \in \mathcal{M}$  so that  $(\bar{a}, y) \in D$ , then  $(\bar{a}, f(\bar{a})) \in D$ . Informally, one says that a Skolem function finds a witness for  $D$ , if there is one. Skolem functions are useful in their own right, both in providing conditions for model completeness, and as a tool used in automated theorem-proving. Any o-minimal model with a group operation (an *o-minimal group*) can also be shown to have definable Skolem functions. The algorithm for determining Skolem functions expands upon the following simple case: if  $D(x, y) \subseteq \mathcal{M}^2$  defines, for every fixed value of  $x$ , an interval, then value of the Skolem function for each  $a$  is the midpoint of the interval defined by the  $D(a, y)$ . O-minimal structures also satisfy the related property of having *uniform elimination of imaginaries*.

*Weakly o-minimal structures* generalize o-minimal structures by allowing each definable subset of the model  $\mathcal{M}$  to be a finite union of *convex sets* which are not necessarily intervals. Consider the ordered group of rational numbers,  $(\mathbb{Q}, +, <)$ . This structure is o-minimal. If we add to the structure the definable set  $P = \{x \in \mathbb{Q} : x < \pi\}$ , then the resulting expansion  $(\mathbb{Q}, +, <, P)$  is not o-minimal: the supremum of the set named by  $P$  is not a rational number, thus  $P$  cannot represent an interval in  $\mathbb{Q}$ . But this set is *convex* in  $\mathbb{Q}$ , and it can be shown that the expanded structure is weakly o-minimal. In fact, it is shown by Baizhanov in [1] that any o-minimal theory, if new convex subsets are introduced, yields a weakly o-minimal theory. A more complex structure of this type is the real-closed valued field  $(R, +, \cdot, 0, 1, <, V)$ , in which  $R$  is a real-closed field with value ring  $V$ . This theory (called *RCVF*) is also weakly o-minimal, and the model theory is explored at length in [3], [4], and [10].

In view of these facts, there is a large program of study concerned with determining which of the properties of o-minimal groups also hold true in the weakly o-minimal case. The authors of [9] distinguish between a *valuational* weakly o-minimal group, in which there is a proper definable subgroup, and a *nonvaluational*

weakly o-minimal group. They showed that while weakly o-minimal groups in general need not have cellular decomposition, the class of nonvaluational weakly o-minimal groups does have an analogue of this property.

## My research projects

My work focuses on more sharply classifying the distinction between valuations and nonvaluational weakly o-minimal structures. In particular, because of the cellular decomposition property enjoyed by nonvaluational weakly o-minimal theories, it is commonly said that a nonvaluational weakly o-minimal theory is as close as possible to an o-minimal theory. My initial results support this conclusion, including a sufficient condition for showing that a model  $\mathcal{M}$  with weakly o-minimal theory is nonvaluational.

**Proposition 1.1.** *Let  $T$  be a weakly o-minimal theory with uniform elimination of imaginaries and definable Skolem functions, and  $\mathcal{M} \models T$ . Then  $\mathcal{M}$  is nonvaluational.*

This proposition actually arises as a simple corollary of a deeper lemma, which may yet have some broader consequences.

**Lemma 1.2.** *Let  $\mathcal{M}$  be a model of a weakly o-minimal theory  $T$  which has definable Skolem functions and uniform elimination of imaginaries. Then there is no equivalence relation  $E$  definable on  $\mathcal{M}$  with infinitely many convex equivalence classes of positive measure.*

It is shown in [10] that under certain limitations, real-closed valued fields have elimination of imaginaries. A possible consequence of this along with the proposition is that such structures will fail to have definable Skolem functions; in future research I hope to be able to determine whether this is the case.

Hoping to understand the implications of the above results, we began studying the class of properly nonvaluational weakly o-minimal models, in order to see whether there in fact are any such models which satisfy the conditions of the proposition. A natural class of these is the nonvaluational weakly o-minimal theories obtained by adding a predicate for a new nonvaluational convex subset to an o-minimal structure. However, this work turned up the surprising result that in fact such structures cannot have definable Skolem functions at all.

**Theorem 1.3.** *Let  $\mathcal{M}$  be an o-minimal expansion of an ordered group in the language  $\mathcal{L}$ . Let  $U$  be a new unary predicate symbol,  $\mathcal{L}' = \mathcal{L} \cup \{U\}$ , and  $\mathcal{M}' = (\mathcal{M}, U)$ , where  $U^{\mathcal{M}'}$  is a downward-closed convex set which defines a properly convex nonvaluational cut. Then  $\mathcal{M}'$  does not have definable Skolem functions in  $\mathcal{L}'$ .*

The proof of this theorem is based in part on work by L. van Den Dries on dense pairs of o-minimal structures (cf. [5]), and relied on my lemma below, which establishes the connection between weakly o-minimal structures and dense pairs of o-minimal structures.

**Lemma 1.4.** *Let  $\mathcal{M}$  be o-minimal with language  $\mathcal{L}$ ; let  $\mathcal{L}' = \mathcal{L} \cup \{U\}$ , and  $\mathcal{M}' = (\mathcal{M}, U)$  with  $U^{\mathcal{M}'}$  a downward-closed nonvaluational convex set, and  $\mathcal{N} = \text{pr}(\mathcal{M} \cup \{b\})$ , where  $b$  realizes  $\text{tp}_{\mathcal{L}}(\text{sup } U/M)$ . Then for any  $X \subseteq M$  definable in  $\mathcal{M}'$ , there is an  $\mathcal{L}$ -formula  $\varphi_X(\bar{x}, y)$  such that  $X = \varphi_X(\mathcal{N}^n, b) \cap \mathcal{M}^n$ .*

Finally, we investigated the question of Skolem functions in valuations models obtained in the same way. In this work, I discovered an algorithmic way to prove the existence of Skolem functions in a subclass of such models.

**Definition 1.5.** Let  $\mathcal{M}$  be an o-minimal expansion of an ordered group, and  $V \subseteq M$  be a convex set. We say that the pair  $(\mathcal{M}, V)$  is *T-immune* if  $V \subsetneq M$  and for any 0-definable function  $F : M \rightarrow M$  and any open convex set  $I \subseteq V^{\mathcal{M}}$ , if  $F \upharpoonright I$  is continuous, then  $F(V) \subseteq V$ .

As an example of *T-immunity*, consider a nonstandard model of the real group,  $\mathcal{M} = (\mathbb{R}^*, +, <, 0)$ , where  $\mathbb{R} \subsetneq \mathbb{R}^*$ , and consider  $V$  interpreted by  $\mathbb{R}$ . Then  $(\mathcal{M}, V)$  is valuations and has a weakly o-minimal theory, and in particular is *T-immune*.

**Theorem 1.6.** *Let  $(\mathcal{M}, +, <, 0, \varepsilon, \dots)$  be an o-minimal expansion of a group with named positive element  $\varepsilon$  in a language  $\mathfrak{L}$  which admits elimination of quantifiers, and  $V \subseteq M$  such that  $(M, V)$  is  $T$ -immune. (Note that since  $\varepsilon \in \mathfrak{L}$ , then  $\varepsilon^M \in V$ .) Let  $c$  be a new constant symbol and  $c^M > 0$  an element of  $M \setminus V$ . Then  $(M, V, c)$  has definable Skolem functions in the language  $\mathfrak{L} \cup \{V, c\}$ .*

L. van den Dries has studied theories with a property known as  $T$ -convexity, which generalizes the notion of  $T$ -immunity. The authors of [7] showed that  $T$ -convex theories also have definable Skolem functions. I am currently working on generalizing the algorithm for calculating Skolem functions in a  $T$ -immune theory to the  $T$ -convex case in order to give an explicit construction.

A natural extension of these results would be a precise set of conditions for definable Skolem functions in any weakly o-minimal theory. For technical reasons, there are many theories which fail to be  $T$ -convex, but may be made so by augmenting the language in a simple way. Modulo a reasonable concept of “almost  $T$ -convexity,” I am investigating now whether it is true that no weakly o-minimal theory obtained by the Baizhanov technique which fails to be “almost  $T$ -convex” has definable Skolem functions. For now, the chief method for doing this is to generalize the concept of dense pairs to theories which may be valuationsal.

## References

- [1] B. Baizhanov, “Expansion of a model of a weakly o-minimal structure by a family of convex predicates,” *J. Symb. Log.* **66** No. 3, 1382-1414 (2001).
- [2] C. Chang, H. J. Keisler, *Model Theory*, Studies in Logic and the Foundations of Mathematics, vol 73 (Elsevier, Amsterdam 1973).
- [3] G. Cherlin, M. A. Dickmann, “Real closed rings I: Residue rings of rings of continuous functions,” *Fund. Math.* **126** 147-183 (1986).
- [4] G. Cherlin, M. A. Dickmann, “Real closed rings II: Model theory,” *Ann. Pure App. Logic* **25** 213-231 (1983).
- [5] L. van den Dries, “Dense pairs of o-minimal structures,” *Fund. Math.* **157**, 61-78 (1998).
- [6] L. van den Dries, *Tame Topology and O-minimal Structures*, London Mathematical Society Lecture Note Series, vol. 248 (Cambridge University Press, Cambridge 1998).
- [7] L. van den Dries, A. H. Lewenberg, “ $T$ -Convexity and tame extensions,” *J. Symb. Log.* **60** No. 1, 74-102 (1995).
- [8] J. Knight, A. Pillay, C. Steinhorn, “Definable sets and ordered structures II,” *Trans. AMS* **295**, 593-605 (1986).
- [9] D. MacPherson, D. Marker, C. Steinhorn, “Weakly o-minimal structures and real closed fields,” *Trans. Amer. Math. Soc.* **352** No. 12, 5435-5483 (2000).
- [10] T. Mellor, “Imaginariness in real closed valued fields,” *Ann. Pure App. Logic* **139**, 230-279 (2006).
- [11] A. Pillay, C. Steinhorn, “Definable sets in ordered structures I,” *Trans. AMS* **295**, 565-592 (1986).
- [12] A. Pillay, C. Steinhorn, “Definable sets in ordered structures III,” *Trans. AMS* **309**, 469-476 (1988).

Christopher S. Shaw  
 Department of Mathematics  
 University of Maryland, College Park  
 schris@umd.edu  
 www.math.umd.edu/~schris